

LIST OF FIGURES

Figure 1-1. Map of the Los Angeles Harbor	3
Figure 3-1. Microbiological sampling stations in Los Angeles Harbor and Cabrillo Beach.....	27
Figure 3-2. Wet- and dry-weather annual geometric means at Los Angeles Harbor surface stations, 2004 and 2005.....	29
Figure 3-3. Annual geometric means for fecal coliform bacteria at Los Angeles Harbor water quality plume tracking stations, 2004 and 2005.....	30
Figure 3-4. Monthly wet- and dry-weather geometric means for indicator bacteria at Cabrillo Beach shoreline stations and rainfall data, 2004.....	31
Figure 3-5. Monthly wet- and dry-weather geometric means for indicator bacteria at Cabrillo Beach shoreline stations and rainfall data, 2005.....	32
Figure 4-1. Water quality sampling stations in Los Angeles Harbor.....	41
Figure 4-2. Calculated salinity anomaly (S_A) values for various dilutions based on mean salinity of 33.346 psu and computed salinities.	43
Figure 4-3a. Temperature and salinity anomaly plots for January, 2004 – May, 2004.....	47
Figure 4-3b. Temperature and salinity anomaly plots for June, 2004 – October, 2004.....	48
Figure 4-3c. Temperature and salinity anomaly plots for November, 2004 – March, 2005.....	49
Figure 4-3d. Temperature and salinity anomaly plots for April, 2005 – August, 2005.....	50
Figure 4-3e. Temperature and salinity anomaly plots for September, 2005 – December, 2005.....	51
Figure 5-1. Sediment chemistry sampling station locations in Los Angeles Harbor.....	58
Figure 5-2. Total metals in Los Angeles Harbor sediment in 2004 and 2005. Concentrations were normalized against fine grain fraction.....	62
Figure 5-3. Total DDT in Los Angeles Harbor sediment in 2004 and 2005. Concentrations were normalized against fine grain fraction.....	62
Figure 5-4. Distribution of % (silt + clay) in Los Angeles Harbor in 2004 and 2005.....	65
Figure 5-5. TOC in Los Angeles Harbor sediment in 2004 and 2005.....	65

Figure 5-6. TOX in Los Angeles Harbor sediment in 2004 and 2005.....	66
Figure 5-7. Trends of copper, nickel, chromium, and zinc at Station HM3 from 2001 to 2005.....	70
Figure 5-8. Trends of copper, nickel, chromium, and zinc at Station HM13 from 2001 to 2005.....	70
Figure 5-9. Trends of mercury, cadmium, arsenic, and lead at Station HM3 from 2001 to 2005.....	71
Figure 5-10. Trends of mercury, cadmium, arsenic, and lead at Station HM13 from 2001 to 2005. Arsenic was not detected in 2002.....	71
Figure 6-1. Macrofaunal sampling stations in Los Angeles Harbor.....	77
Figure 6-2. Parsimony analysis of endemicity (PAE) cladogram for the 2004-2005 biennial stations.....	85
Figure 6-3. Non-metric multi-dimensional scaling (NMDS) using patristic distances derived from the PAE cladogram for the 2004-2005 biennial stations. Stations (a), season (b), BRI response level (c), and total taxonomic distinctness (d) are depicted as factors overlaid on the same NMDS configuration in different perspectives.....	87
Figure 6-4. Parsimony analysis of co-occurring species (PACOS) cladogram showing the taxa relationships based on where they occurred during the 2004-2005 biennium.....	88
Figure 6-5. Parsimony analysis of endemicity (PAE) cladogram of Summer 2004 and 2005 sampling events. The outgroup is the basal outside-breakwater pair.....	89
Figure 6-6. Non-metric multi-dimensional scaling (NMDS) of Summer 2004 and Summer 2005 stations from patristic distances derived from the PAE cladogram with mapping of stations (a), zinc (b), NERL (c), ERLq range (d), toxicity range (e), and sand percent range (f). Red dashed circles represent outside-breakwater Station HM13. Outfall events (HM3) are identified by red arrows.....	91
Figure 6-7. Principal components analysis (PCA) of the PAE cladogram with sediment granulometry (clay, silt, and sand) for summer 2004 and 2005. Independent mapping of silt (left plot) and sand (right plot) back onto the ordination shows the relative distributions of each.....	92
Figure 7-1. Trawl sampling stations in Los Angeles Harbor. Abandoned or former station positions are depicted in gray.....	103
Figure 7-2. <i>Genyonemus lineatus</i> from Miller and Lea (1972)......	109

Figure 7-3. Size-class (cm) distribution of <i>Genyonemus lineatus</i> during the 2004-2005 biennium.....	109
Figure 7-4. Size-class (cm) distribution of <i>Seriphus politus</i> during the 2004-2005 biennium.....	110
Figure 7-5. <i>Crangon nigromaculata</i> drawing.....	111
Figure 7-6. <i>Philine auriformis</i> drawing.....	111
Figure 7-7. Parsimony analysis of endemicity (PAE) q-mode cladogram showing the sampling event relationships based on the combined fish and mega invertebrate taxa inventories of the 2004 – 2005 biennium.....	118
Figure 7-8. Non-metric multi-dimensional scaling (NMDS) of patristic distances derived from the parsimony analysis of endemicity (PAE) cladogram showing the sampling event relationships based on the combined fish and megainvertebrate taxa inventories of the 2004-2005 biennium. Three treatment levels of a location factor are employed.....	119
Figure 7-9. Parsimony analysis of co-occurring species (PACOS) cladogram showing the associations of combined fish and megainvertebrate species based on where (sampling events) they occurred during the 2004-2005 biennium. The paired, condensed views of the same cladogram offer species abundance mapping that correspond directly with clade features (green ellipse) of the larger tree view.	120
Figure 8-1. Trawl stations in Los Angeles Harbor during 2004 and 2005 survey.....	126
Figure 8-2. Temporal trend of selected pollutants and % lipid in muscle tissue of white croaker at the TITP outfall (HT7) from 2000 – 2005. Arsenic was not detected in 2001 and 2002.	133
Figure 8-3. Concentrations of selected pollutants in muscle and liver tissues near the TITP outfall during the 2004 and 2005 sampling seasons.	134

LIST OF TABLES

Table 1-1.	Summary of the Los Angeles Harbor Monitoring Programs, including the 1993 NPDES requirements, interim program, the Pier 400 monitoring program, and the new 2005 NPDES requirements.....	4
Table 2-1a.	Constituents measured in the effluent monitoring program from January 2004 to May 2005.	11
Table 2-1b.	Constituents measured in the effluent monitoring program from May 2005 to December 2005.	12
Table 2-2a.	The NPDES effluent limits and annual averages of major effluent constituents in 2004	15
Table 2-2b.	The NPDES effluent limits and annual averages of major effluent constituents in 2005.	15
Table 2-3a.	2004 monthly averages for major wastewater constituents.....	16
Table 2-3b.	2005 monthly averages for major wastewater constituents.....	16
Table 2-4.	Annual average of major wastewater constituents from 1995 to 2005.....	17
Table 2-5a.	The NPDES effluent limits and annual averages of priority pollutant inorganics in 2004.....	18
Table 2-5b.	The NPDES effluent limits and annual averages of priority pollutant inorganics in 2005.....	18
Table 2-6.	Annual average concentrations of metal priority pollutants in influent and effluent from 1995 to 2005.....	19
Table 2-7a.	The NPDES effluent limits and annual averages of priority pollutant organics in 2004.....	20
Table 2-7b.	The NPDES effluent limits and annual averages of priority pollutant organics in 2005.....	21
Table 3-1.	Summary of the Los Angeles Harbor and Cabrillo Beach Microbiological Monitoring Program for 2004-2005.....	26
Table 3-2.	Number of dry-weather * non-compliance days and percent compliance for NPDES Bathing Water and Shellfish Harvesting coliform limits at Cabrillo Beach shoreline stations in 2004.....	33

Table 3-3.	Number of dry-weather * non-compliance days and percent compliance for NPDES Bathing Water and Shellfish Harvesting coliform limits at Cabrillo Beach shoreline stations from January to May 2005.	33
Table 3-4.	Number of dry-weather * non-compliance days and percent compliance for NPDES Bathing Water and Shellfish Harvesting coliform and enterococcus limits at Cabrillo Beach shoreline stations from June to December 2005	34
Table 3-5.	Number of occurrences and percent compliance with all materials of sewage origin at Cabrillo Beach shoreline stations in 2004-2005.	34
Table 4-1.	NPDES receiving water limitations for the Terminal Island Treatment Plant.....	40
Table 4-2.	Minimum and maximum values from temperature (°C), salinity (psi), dissolved oxygen (mg/L), and pH measured during each water quality survey during 2004-2005	45
Table 4-3.	Summary of 2004-2005 survey dates, oceanic conditions, tides, nearfield and farfield station locations, and direction. Sampling was conducted between 0800 and 1500.	46
Table 4-4.	Summary of acute toxicity results.....	52
Table 4-5.	Summary of chronic toxicity results using <i>Haliotis rufescens</i>	53
Table 4-6.	Probability estimates, with confidence limits, of wastewater field occurrence at station and depth calculated from salinity anomaly values >0.57 measured July 1997 – December 2005	54
Table 5-1.	Grain size distribution of sediments at Los Angeles Harbor sites surveyed in summer 2004 and 2005	60
Table 5-2.	Cyanide, dissolved sulfides, TOC, and TOX in Los Angeles Harbor sediment in summer 2004 and 2005. Concentrations were normalized against fine grain fraction.	61
Table 5-3.	Metals detected in sediments at stations HM2 - HM4 and HM6 - HM13 of Los Angeles Harbor in summer 2004 and 2005. Concentrations were normalized against fine grain fraction.....	61
Table 5-4.	BNA and pesticides detected in sediments at stations HM2 - HM4 and HM6 - HM13 of Los Angeles Harbor in summer 2004 and 2005. Concentrations were normalized against fine grain fraction.....	63
Table 5-5.	Metals in sediments of Los Angeles Harbor in summer 2004 and 2005, and at a reference site surveyed from 1984 – 1987. Concentrations were normalized against fine grain fraction.....	66

Table 5-6.	BNA and pesticides detected in sediments at stations of Los Angeles Harbor in summer 2004 and 2005. Concentrations were normalized against fine grain fraction	67
Table 5-7.	Concentrations of metals and organic pollutants in Los Angeles Harbor, summer 2004 and 2005 with comparison to concentrations in Southern California Bight (SCB) in 1998	67
Table 5-8.	Comparison of contaminant levels at TITP outfall, stations in Santa Monica Bay during summer 2004, 2005, and a reference site. Concentrations were normalized against fine grain fraction.....	69
Table 5-9.	Temporal pattern of total DDT and metals concentrations at HM13 (outside breakwater of the Harbor) and HM3 (outfall). Concentrations were normalized against fine grain fraction.....	69
Table 5-10.	Concentrations of metals and organic pollutants in sediments of Los Angeles Harbor, summer 2004 and 2005, with comparison to Effective Levels* (ER-L and ER-M). Concentrations were normalized against fine grain fraction.....	72
Table 6-1.	Los Angeles Harbor 2004-2005 biennium macrofaunal community parameters, various biodiversity measures, and phylodiversity indices by replicate (sample) including species (taxa) richness (S), abundance (N), Margalef's species richness (d), Pielou's evenness (J') Shannon-Weiner function (H' log base e), Simpson's index (1- λ'), taxonomic distinctness (Δ^*), variation in taxonomic distinctness (Λ^+), total taxonomic distinctness (s Δ^+), and total phylogenetic diversity (s Φ^+), and benthic response index (BRI).....	82
Table 6-2.	Traditional diversity indices Pearson Product Moment correlation coefficients and P-values (in parentheses) pair-wise matrix.....	83
Table 6-3.	Phylodiversity indices versus taxa richness Pearson Product Moment correlation coefficients and P-values (in parentheses).....	84
Table 6-4.	Distribution of effects range-low (ERL) value exceedances (gray shading) for Summer 2004 and Summer 2005 surveys. Boldface values depict exceedances. Gray font depicts values below ERL.....	90
Table 7-1.	Los Angeles Harbor 2004-2005 biennium fish and megainvertebrate community parameters, various biodiversity measures, and phylodiversity indices by replicate (sample) including species (taxa) richness (S), abundance (N), Margalef's species richness (d), Pielou's evenness (J') Shannon-Weiner function (H' log base e), Simpson's index (1- λ'), taxonomic distinctness (Δ^*), variation in taxonomic distinctness (Λ^+), total taxonomic distinctness (s Δ^+), and total phylogenetic diversity (s Φ^+), and fish response index (FRI).	106
Table 7-2.	Fish species abundances, total abundances per event, total abundances per species, and number of species per event in Los Angeles Harbor, 2004	107

Table 7-3.	Fish species abundances, total abundances per event, total abundances per species, and number of species per event in Los Angeles Harbor, 2005	108
Table 7-4.	Los Angeles Harbor 2004-2005 biennium megainvertebrate community parameters, various biodiversity measures, and phylodiversity indices by replicate (sample) including species (taxa) richness (S), abundance (N), Margalef's species richness (d), Pielou's evenness (J') Shannon-Weiner function (H' log base e), Simpson's index ($1 - \lambda'$), taxonomic distinctness (Δ^*), variation in taxonomic distinctness (Δ^+), total taxonomic distinctness ($s\Delta^+$), and total phylogenetic diversity ($s\Phi^+$)	112
Table 7-5.	Megainvertebrate taxa abundances, total abundances per event, total abundances per taxon, and number of taxa per event in Los Angeles harbor, 2004	113
Table 7-6.	Megainvertebrate taxa abundances, total abundances per event, total abundances per taxon, and number of taxa per event in Los Angeles harbor, 2005	114
Table 7-7.	Taxa richness and biomass for Los Angeles Harbor, 2004-2005 biennium	116
Table 7-8.	Combined fish and megainvertebrate community parameters, 2004-2005 biennium. Indices and abbreviations follow those of Table 7-1	117
Table 8-1.	Location and number of trawl and sport fish samples collected during 2004 and 2005.	127
Table 8-2.	Metals, cyanide, and organics found in muscle tissues of white croaker at HT7 in Los Angeles Harbor during 2004 to 2005 period. Bold numbers indicate DDT and PCB levels that exceed the 0.1 mg/kg limit set by OEHHA (1991)	128
Table 8-3.	Metals, cyanide, and organics found in muscle tissues of Queenfish at HT12 and liver tissues of white croaker at HT7 in Los Angeles Harbor during 2005 period. Bold numbers indicate DDT and PCB levels that exceed the 0.1 mg/kg limit set by OEHHA (1991).	129
Table 8-4.	Selected metals and organics found in muscle tissues of white croaker in Los Angeles Harbor and Santa Monica Bay during 2005. Bold numbers indicate DDT and PCB levels that exceed the 0.1 mg/kg limit set by OEHHA (1991).	131
Table 8-5.	Concentrations of pollutants in muscle tissues of white croaker and sediments near the TITP outfall stations, HT7 and HM3, respectively, during the 2004 and 2005 sampling seasons	132
Table C-1.	Method detection limits of metals in sediments and tissues	C6